

Soybean Row Spacing



on-farm network
PARTICIPATORY • PRECISE • PROACTIVE

Do soybeans on narrow spacing out-perform soybeans on wide spacing? Find out!

Suggested Treatment Comparisons

- Wide row at reduced seeding rate vs. narrow row at normal seeding rate

OR

- Wide row vs. narrow row (consistent seeding rate)

Trial Design

- Follow a randomized layout
- Minimum four replicates
- Strips need to be at least as wide as one combine pass
- Strips should run the length of the field, excluding the headlands

Your Fields, Your Results

- Established with your equipment
- Under your management practices
- Harvested with your equipment
- MPSG support at establishment & harvest
- MPSG collects data throughout the season
- Results packaged together for you in a short report
- Must be a member in good standing

Headland

Wide	Narrow	Wide	Narrow	Narrow	Wide	Narrow	Wide	Wide	Narrow	Wide	Narrow
1	2	3	4	5	6	7	8	9	10	11	12

Headland

Choose the treatment combination that works best for your farm!

Decision Making, Simplified
Your trial includes:

- ✓ Plant stand data
- ✓ Canopy closure data
- ✓ Weed pressure data
- ✓ Yield results
- ✓ Economics analysis

Your data, working for your farm

MANITOBA
Pulse Soybean
GROWERS

To participate in the trials, contact

Chris Forsythe 204.751.0439 chris@manitobapulse.ca

or sign up at **manitobapulse.ca/on-farm-network**