Soybean Inoculant

Single vs. None or Double vs. Single

on-farm network
PARTICIPATORY • PRECISE • PROACTIVE

Does a reduction in soybean inoculant affect nodulation and yield? Find out!

Suggested Treatment Comparisons

- ❑ Double inoculant vs. single inoculant

OR

- ❑ Single inoculant vs. no inoculant

Headland

Double (or none)	Single	Double (or none)	Single	Single	Double (or none)	Single	Double (or none)	Double (or none)	Single	Double (or none)	Single
1	2	3	4	5	6	7	8	9	10	11	12

Headland

***3-year soybean history required for single vs. none**

***2-year soybean history required for double vs. single**

Trial Design

- Follow a randomized layout
- Minimum four replicates
- Strips need to be at least as wide as one combine pass
- Strips should run the length of the field, excluding the headlands

Your Fields, Your Results

- Established with your equipment
- Under your management practices
- Harvested with your equipment
- MPSG support at establishment & harvest
- MPSG collects data throughout the season
- Results packaged together for you in a short report
- Must be a member in good standing

Decision Making, Simplified
Your trial includes:

- ✓ **Plant stand data**
- ✓ **Nodulation ratings**
- ✓ **Yield results**
- ✓ **Economics analysis**

Your data, working for your farm

MANITOBA Pulse Soybean GROWERS

To participate in the trials, contact

Megan Bourns • 204.751.0439 • megan@manitobapulse.ca

or sign up at manitobapulse.ca/on-farm-network