

# Pea Inoculant

*Single vs. None or Double vs. Single*



**on-farm network**  
PARTICIPATORY • PRECISE • PROACTIVE

***Does a change in pea inoculation strategy lead to a change in nodulation and yield? Find out!***

## Suggested Treatment Comparisons

- Double inoculant vs. single inoculant

OR

- Single inoculant vs. no inoculant

## Trial Design

- Follow a randomized layout
- Minimum four replicates
- Strips need to be at least as wide as one combine pass
- Strips should run the length of the field, excluding the headlands

Headland

Double (or none)	Single	Double (or none)	Single	Single	Double (or none)	Single	Double (or none)	Double (or none)	Single	Double (or none)	Single
1	2	3	4	5	6	7	8	9	10	11	12

Headland

***Test the strategies that make the most sense for your farm!***

## Your Fields, Your Results

- Established with your equipment
- Under your management practices
- Harvested with your equipment
- MPSG support at establishment & harvest
- MPSG collects data throughout the season
- Results packaged together for you in a short report
- Must be a member in good standing

***Decision Making, Simplified***  
***Your trial includes:***

- ✓ Plant stand data
- ✓ Nodulation ratings
- ✓ Yield results
- ✓ Economics analysis

***Your data, working for your farm***



To participate in the trials, contact

**Leanne Koroscil • 204.751.0439 • [leanne@manitobapulse.ca](mailto:leanne@manitobapulse.ca)**

**or sign up at [manitobapulse.ca/on-farm-network](http://manitobapulse.ca/on-farm-network)**